


Black-Eared Wheatear

Oenanthe hispanica



SEASONAL BEHAVIOUR

On passage seen from March to May, and from August to early October.

DISTRIBUTION

In Huelva nests in roadside embankments in *Cistus* scrub and some mine tailings, as in Sevilla and Córdoba. In the latter province there are some pairs on crests and ridges (Guadiato valley). In Jaén breeds on abandoned mines (Los Guindos) and well developed crags (El Puntal, Selladores, etc).

THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	NT	-

WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			

STATUS

Migrant and summer visitor.

ABUNDANCE

Scarce.

HABITAT

On passage seen in the same type of places as the former species. Breeds in two slightly different habitats: rocky outcrops and crags on the one hand and, on the other, roadside embankments, mine tailings, edges of *Cistus* scrub and hilltops with stony soil and scant vegetation.