


Black-Headed Gull

Larus ridibundus



SEASONAL BEHAVIOUR

Seen for much of the year, since the early birds arrive in July and the last leave in March or even April.

DISTRIBUTION

Appears on all sizeable reservoirs though never in high numbers. Also seen on the Guadalquivir and the lower reaches of the Guadiato.

THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	-

WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			

STATUS

Winter visitor and migrant.

ABUNDANCE

Scarce, with a slightly upward trend in recent years.

HABITAT

Mainly reservoirs but also smaller bodies of open water and wide stretches of the biggest rivers.