



Black Tern

Chlidonias niger



SEASONAL BEHAVIOUR

In theory it should turn up on both spring and autumn passage but the only actual records refer to the latter.

DISTRIBUTION

Has been seen in Huelva (Aracena Reservoir) and Córdoba (Puente Nuevo Reservoir).

THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
EN	EN	EN	CR

WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			

STATUS

Migrant.

ABUNDANCE

Rare.

HABITAT

Big reservoirs.