

## Bonelli's Eagle

*Aquila fasciata*




### THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
VU	VU	EN	VU

### WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			



### SEASONAL BEHAVIOUR

Seen all year.

### DISTRIBUTION

Token presence only in Huelva and Jaén (one pair in each); the bulk of the population lives in Sierra Morena Sevillana and roundabout (15 pairs) and in the Córdoba portion of Sierra Morena (16 pairs). Favoured areas for natal dispersal are Chanza valley, the foothills of Montoro and the plains of Guadalcanal.

### STATUS

Sedentary, with some vagrants.

### ABUNDANCE

Uncommon in the central band and rare elsewhere (33 pairs).

### HABITAT

Favoured habitat is Mediterranean woodland although it also shows in olive-growing areas if there are nearby cliffs and rocky outcrops. As a species nests mainly on cliff ledges though there are many tree-nesting pairs in Sierra Morena.