


Common Sandpiper

Actitis hypoleucos



SEASONAL BEHAVIOUR

Like the former species has a very long migration period and winter visitors come back early; birds are also present in spring and summer so there are records in every month of the year.

DISTRIBUTION

Distributed throughout the whole territory.

COMMENTS

There is a record of two birds with suspected breeding behaviour on River Sotillo (on the border between Seville and Badajoz) in May 2009, though breeding was not eventually proven. It has bred on rivers and reservoirs of southern Huelva in recent years.

THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	-

WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			

STATUS

Migrant and winter visitor, with some non-breeding summering birds.

ABUNDANCE

Scarce.

HABITAT

Fonder of the shores of large reservoirs than the Green Sandpiper, though also seen on rocky rivers.