


## Feral Pigeon

*Columba livia* var. *domestica*



### SEASONAL BEHAVIOUR

Seen all year.

### DISTRIBUTION

Distributed throughout the whole territory.

### THREATS

There are unlikely now to be any colonies of pure Rock Doves left in Sierra Morena, reflecting the dire situation of this species in Spain due to the genetic contamination of *Columba livia* after centuries of cross-breeding with feral pigeons.

### THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	-

### WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			

### STATUS

Sedentary.

### ABUNDANCE

Common.

### HABITAT

Breeds almost exclusively on human constructions both in towns and open country (bridges, farmsteads, reservoir dams, hermitages, etc) but there are some cliffnesting colonies (Peña de Arias Montano, Cerro del Hierro, Sierra de Hornachuelos, Los Órganos de Despeñaperros, etc).