


## Quail

*Coturnix coturnix*



### SEASONAL BEHAVIOUR

The bulk arrive in March and leave in August-September, through there are records in every month of the year.

### DISTRIBUTION

Most frequent in the campiña of Guadalquivir valley, though could easily be seen in any of the four provinces. Overwintering birds have been recorded in Huelva and Córdoba.

### THREATS

One of the species most affected by the declining farming activity in Sierra Morena. In the past, when small plots were sown up in valleys and on hillsides, it was commoner and more widely distributed.

### THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	DD	-

### WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
<b>H-1</b>	<b>SE-1</b>	<b>CO-1</b>	<b>J-1</b>
H-2	<b>SE-2</b>	CO-2	J-2
H-3	SE-3	<b>CO-3</b>	<b>J-3</b>
H-4	<b>SE-4</b>	<b>CO-4</b>	J-4
<b>H-5</b>	<b>SE-5</b>	CO-5	J-5
<b>H-6</b>	<b>SE-6</b>	CO-6	J-6
H-7	SE-7	CO-7	<b>J-7</b>
H-8	<b>SE-8</b>	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	<b>SE-10</b>	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	<b>J-13</b>
<b>H-14</b>	<b>SE-14</b>	CO-14	J-14
H-15	SE-15	<b>CO-15</b>	J-15
H-16			
H-17			

### STATUS

Summer visitor with a few winter records.

### ABUNDANCE

Scarce in general though relatively numerous in some parts.

### HABITAT

Pastureland, grassland and cropland. Sometimes in clearings, dehesas and hilltops.