

Red-Necked Nightjar

Caprimulgus ruficollis



THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	DD

WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17	[






























Summer visitor and migrant.

ABUNDANCE

Uncommon in general.

HABITAT

Sunny areas with open spaces and some shrub cover and/or clumps of trees, like patches of scrub, clearings in pinewoods and Eucalyptus woods, cropland, pastureland, valleys of rivers and streams,



SEASONAL BEHAVIOUR

Most stay with us from April to September.

DISTRIBUTION

Absent or scarce in most of Huelva and Seville. More numerous in Córdoba and Jaén.

COMMENTS

Its percussive song has won it onomatopoeic local nicknames like capacho, gapacho and especially zumaya, zamaya or azamaya. Sometimes enters towns to hunt moths round streetlights.









