


## Sedge Warbler

*Acrocephalus schoenobaenus*



### SEASONAL BEHAVIOUR

Detected above all on spring passage (late February-April) when the males are singing.

### DISTRIBUTION

Can be found in suitable areas of all four provinces.

### THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	-

### WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	<b>SE-1</b>	CO-1	J-1
H-2	<b>SE-2</b>	CO-2	J-2
H-3	SE-3	<b>CO-3</b>	J-3
H-4	SE-4	CO-4	<b>J-4</b>
H-5	SE-5	CO-5	J-5
H-6	SE-6	<b>CO-6</b>	J-6
H-7	SE-7	<b>CO-7</b>	J-7
H-8	SE-8	CO-8	J-8
<b>H-9</b>	<b>SE-9</b>	CO-9	<b>J-9</b>
<b>H-10</b>	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	<b>CO-12</b>	J-12
H-13	<b>SE-13</b>	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	<b>CO-15</b>	<b>J-15</b>
H-16			
<b>H-17</b>			

### STATUS

Migrant.

### ABUNDANCE

Scarce.

### HABITAT

Drops into riparian thickets of Oleander and Tamujo and wetland vegetation. Sometimes also the shadier scrubby areas.