

Starling

Sturnus vulgaris



SEASONAL BEHAVIOUR

Likeliest to be seen from October to March.

DISTRIBUTION

Might be found in any suitable habitat throughout the region though in general it is scarce and tricky to see.

COMMENTS

Joins flocks of the much more abundant Spotless Starling, which it greatly resembles in winter plumage.

THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	-

WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			

STATUS

Winter visitor.

ABUNDANCE

Scarce.

HABITAT

Cropland in the campiña, dehesas and upland livestock pastures.