


# Wheatear

*Oenanthe oenanthe*



### SEASONAL BEHAVIOUR

Much more frequent on post-breeding passage (especially August to October) than in the spring.

### DISTRIBUTION

The whole territory.

### THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	-

### WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
<b>H-1</b>	<b>SE-1</b>	<b>CO-1</b>	<b>J-1</b>
H-2	<b>SE-2</b>	<b>CO-2</b>	J-2
H-3	SE-3	CO-3	<b>J-3</b>
H-4	<b>SE-4</b>	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
<b>H-8</b>	SE-8	<b>CO-8</b>	J-8
H-9	SE-9	CO-9	J-9
H-10	<b>SE-10</b>	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
<b>H-13</b>	SE-13	CO-13	<b>J-13</b>
<b>H-14</b>	<b>SE-14</b>	CO-14	J-14
H-15	SE-15	<b>CO-15</b>	J-15
H-16			
H-17			

### STATUS

Migrant.

### ABUNDANCE

Uncommon.

### HABITAT

Turns up in open areas free of vegetation, like pastureland, cropland and *campiña*, patches of low scrub, etc, and also on crags, mines and in rides and firebreaks, etc.