


Whitethroat

Sylvia communis



SEASONAL BEHAVIOUR

Recorded on both pre- and post-breeding passage though more common in the latter, especially in August and September.

DISTRIBUTION

Seen in the whole cordillera.

THREAT CATEGORY

CNEA	CAEA	LRAE	LRVA
-	-	-	NT

WHERE TO SEE IT

HUELVA	SEVILLA	CÓRDOBA	JAÉN
H-1	SE-1	CO-1	J-1
H-2	SE-2	CO-2	J-2
H-3	SE-3	CO-3	J-3
H-4	SE-4	CO-4	J-4
H-5	SE-5	CO-5	J-5
H-6	SE-6	CO-6	J-6
H-7	SE-7	CO-7	J-7
H-8	SE-8	CO-8	J-8
H-9	SE-9	CO-9	J-9
H-10	SE-10	CO-10	J-10
H-11	SE-11	CO-11	J-11
H-12	SE-12	CO-12	J-12
H-13	SE-13	CO-13	J-13
H-14	SE-14	CO-14	J-14
H-15	SE-15	CO-15	J-15
H-16			
H-17			

STATUS

Migrant.

ABUNDANCE

Common.

HABITAT

Turns up in patches of scrub, orchards, hedgerows, *dehesas*, olive groves, riversides, etc, without any strict preference for either. In summer sticks close to sources of water.